



PATENT ABSTRACTS OF JAPAN

(11) Publication number: 09048770 A
 (43) Date of publication of application: 18.02.1997

(51) Int. Cl. C07D263/44

A61K 31/41,	A61K 31/42,	A61K 31/42,	A61K 31/425,	A61K 31/425,
A61K 31/425,	A61K 31/425,	A61K 31/425,	A61K 31/44,	A61K 31/47,
A61K 31/495,	C07D271/06,	C07D277/34,	C07D413/12,	C07D413/12,
C07D413/12,	C07D413/12,	C07D413/14,	C07D413/14,	C07D417/12,
C07D417/12,	C07D417/12,	C07D417/12,	C07D417/12,	C07D417/12,
C07D417/12,	C07D417/12,	C07D417/12,	C07D417/14,	C07D417/14,
C07D417/14,	C07D417/14,	C07D417/14,	C07D417/14,	C07D417/14,
C07D493/04,	C07D493/04,			

(21) Application number: 08136609
 (22) Date of filing: 30.05.1996
 (30) Priority: 02.06.1995 JP 07137002

(71) Applicant: SANKYO CO LTD
 (72) Inventor: YANAGISAWA HIROAKI
 FUJITA TAKESHI
 FUJIMOTO KOICHI
 WADA KUNIO
 OGUCHI MINORU
 YOSHIOKA TAKAO
 FUJIWARA TOSHIHIKO
 HORIKOSHI HIROYOSHI

(54) AROMATIC OXYIMINO DERIVATIVE


lowed by deprotecting the resultant compound.

COPYRIGHT: (C) 1997, JPO


(57) Abstract:

PROBLEM TO BE SOLVED: To obtain an aromatic oxyimino derivative having remedying effect on insulin resistance, hyperlipemia, hyperglycemia, obesity, etc., excellent in safety, thus useful for preventing and treating these diseases.

SOLUTION: This new compound (salt) is expressed by formula I [R^1 is H, a 1-6C alkyl, 1-4C alkoxy, 1-4C alkythio, etc.; R^2 is a 1-6C alkylene; A is methylene, O, S, etc.; B is (1-3 R^1 (except H)-substituted) 5- or 6-membered heterocyclic divalent group, etc.; X is (2,4,5-trihydro-3H-2,4-dioxo-thiazol-5-yl) methyl, etc.; Y is S, O, etc., e.g. 5-(4-{2-[(6,7-dichloroindan-1-yl) iminoxy]ethoxy}benzyl)-2,4-thiazolidindione. The compound of formula I is obtained by condensation reaction between a compound of formula II (U is OH, halogen, etc.) and a compound of formula III (X' is a group resulted by protecting the NH group at X) fol-



I



II



III